

altoCIRRUS[®]

Fogging Concentrate

Specimen Label

- For use in homes, apartments, food and non-food areas of: food processing plants, warehouses, office buildings, hotels, motels, schools, theatres, hospitals and other areas listed
- Designed for use through ULV and fogging equipment
- Delivers fast knockdown and kill of listed insects

ACTIVE INGREDIENTS:

* (RS)-2-Methyl-4-oxo-3-(2-propynyl) cyclopent-2-enyl-(1RS)-cis, trans-chrysanthamate	0.67%
** Piperonyl butoxide, Technical	2.00%
*** N-Octyl bicycloheptene dicarboximide.....	3.00%
**** OTHER INGREDIENTS	94.33%
	100.00%

* ETOC[®]

** Equivalent to 1.60% (butylcarbityl) (6-propylpiperonyl) ether and 0.40% related compounds

*** MGK[®] 264, Insecticide Synergist

**** Contains petroleum distillate



KEEP OUT OF REACH OF CHILDREN

CAUTION

See inside for first aid and precautionary statements.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ALL DIRECTIONS COMPLETELY BEFORE USE

GENERAL INFORMATION

Remove pets and birds and cover fish aquariums before spraying.

Do not allow children or pets to contact treated surfaces until spray has dried.

GENERAL: This concentrate is designed for use undiluted through mechanical aerosol or Ultra Low Volume (ULV) generators, as well as thermal and conventional fogging or spraying equipment.

DILUTION: It can be diluted with a white mineral oil conforming to regulation 21 CFR 178.3620(b), or an odorless light petroleum hydrocarbon conforming to 21 CFR 172.882 or 40 CFR 180.1001(c) & (e). Combine oil with this concentrate and mix well just prior to application.

TANK-MIXING: This concentrate may be tank mixed with NyGuard[®] Insect Growth Regulator, or other appropriately labeled insect control products, unless prohibited by the other product's label. When using a tank-mix, always make applications following the most restrictive label language of the products being mixed.

NOTE: If there is a need to treat such areas as conduits, motor housings, junction and switch boxes or other electrical equipment, de-energize electrical equipment prior to application. After application, allow time for oil to evaporate before starting motors or placing switches in the "ON" position.

RESTRICTIONS

Product cannot be used: 1) in areas where animals raised for milk and meat (human food animals) are present; 2) in any areas where animal feeds are prepared, processed, or stored, and; 3) in any truck beds or truck trailers, freight/tank/container, railroad cars, ship holds, or ship and air freight containers that are used to transport unbagged (bulk) feeds.

Do not apply to classrooms or patient rooms in hospitals and nursing homes while occupied.

Do not apply to a given area more than once every 4 days.

FOOD HANDLING ESTABLISHMENT APPLICATIONS

Do not apply directly to food. Do not apply as a Space Spray or general surface treatment while food processing is underway. Careful spot and/or crack and crevice application may be used while the facility is in operation provided exposed food is covered or removed from the area being treated prior to application. Foods should be removed or covered during treatment. All food processing surfaces and equipment should be covered or washed with an effective cleaning compound followed by a potable water rinse. In the home, all food processing surfaces, dishes and utensils should be covered during treatment, or thoroughly washed before use.

Insects controlled:

Almond Moths	Cereal Beetles	Coffee Bean Weevils	Earwigs	Hide Beetles	Mosquitoes	Small Flying Moths	Waterbugs
Angoumois Grain Moths	Cheese Mites	Weevils	Face Flies	Horn Flies	Mushroom Flies	Spider Beetles	Weevils
Ants	Cheese Skippers	Confused Flour Beetles	Firebrats	Hornets	Red Flour Beetles	Spiders	Yellow Jackets
Bed Bugs	Chocolate Crickets	Dark Mealworms	Fleas	House Flies	Rice Weevils	Stable Flies	Yellow Mealworms
Blister Beetles	Moths	Dark Mealworms	Flies	Indian Meal Moths	Rusty Grain Beetles	Stink Bugs	Ticks
Blood Sucking Lice	Cigarette Beetles	Darkling Beetles	Fruit Flies	Khapra Beetles	Saw-toothed Grain Beetles	Tobacco Moths	Vinegar Flies
Boxelder Bugs	Clothes Moths	Dried Fruit Beetles	Gnats	Leaf Hoppers	Scorpions	Warehouse Beetles	Wasps
Cadelles	Clover Mites	Drugstore Beetles	Grain Mites	Lesser Grain Borers	Silverfish		
Carpet Beetles	Cockroaches	Harlequin Bugs	Granary Weevils	Mediterranean Flour Moths			
Centipedes							

For use in and around:

Apartments	Cafeterias	Diners	Garbage Compactors	Jails	Museums	Restaurants	Tobacco Warehouses
Arcades	Campers	Dining Halls	Garbage Trucks	Kennels	Mushroom Processing	Retail Stores	Tool Sheds
Auditoriums	Canneries	Dog Houses	Gas Stations	Kitchens	Nursing Homes	Rice Mills	Trains
Bakeries	Cargo Aircraft	Dormitories	Grain Elevators	Laboratories	Office Buildings	Schools	Trucks
Banquet Facilities	Carpets	Drug Stores	Grain Harvesting & Handling Equipment	Laundromats	Office Buildings	Smoke Shops	Tunnels
Barns	Churches	Dumpsters	Grain	Lavatories	Parking Garages	Stables	Tunnels
Bars	Circus Tents	Egg Processing Plants	Harvesting & Handling Equipment	Libraries	Passenger Railroad Cars	Storage Bins	Universities
Basements	City Missions	Ethanol Plants	Equipment	Liquor Distilleries	Pool Halls	Storages	Utilities
Bathrooms	Closets	Factories	Grain Mills	Liquor Stores	Poultry Housing	Facilities	Utility Rooms
Beverage Plants	Communication Centers	Food Service Storage and Preparation Areas	Granaries	Locker Rooms	Prisons	Stored Food in Multi-walled Packaging or Cloth Bags	Utility Rooms
Boats	Concession Stands	Freight Containers	Grocery Stores	Marina Buildings	Public Housing	Warehouses	Voids (i.e. attics, false ceilings)
Bottling Plants	Condominiums	Fruit Packing Sheds	Healthcare Facilities	Mausoleums	Public Storages	Stores	Walls & Equipment
Breweries	Convenience Stores	Garages	Homes (households)	Meat Packing Plants	Rabbit Processing Plants	Supermarkets	Warehouses
Brewery Warehouses	Court Houses	Garbage Bins	Hospices	Mills	Railroad Cars	Textile Mills & Warehouses	Wholesale Stores
Buildings	Day Care Centers	Garbage Cans	Hospitals	Mobile Homes	Recreational Vehicles (RV's)	Theatres	Wine Cellars
Busses	Department Stores		Hotels	Morgues			Wineries
Cabinets			Industrial Installations	Motels	Recycling Facilities		Zoos
Cabins			Institutions	Movie Theatres			

UNDILUTED APPLICATIONS

SPACE SPRAY APPLICATION

TO KILL FLYING INSECTS LISTED ABOVE: Close doors, windows and shut off all ventilating equipment. Then with a mechanical aerosol generator (no droplets over 50 microns in diameter and 80% less than 30 microns) apply undiluted concentrate at a dosage rate of approximately 1 fl. oz./1,000 ft.³ of room space. Direct spray toward upper areas of room. Do not remain in treated areas after application, and leave room closed for at least 1 hour. Ventilate thoroughly before occupants are allowed to re-enter.

TO KILL CRAWLING INSECTS LISTED ABOVE: Close doors, windows and shut off all ventilating equipment before application. Use concentrate undiluted and apply through equipment which can deliver a pin stream spray. Direct a pin stream spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects harbor. Contact as many insects as possible with the spray. Do not allow spray to contact stored food. Then, with a mechanical aerosol generator (no droplets over 50 microns in diameter and 80% less than 30 microns) apply undiluted product at the rate of 1 oz./1,000 ft.³ of space. Leave room closed for at least 1 hour after application. Do not remain in treated areas and ventilate thoroughly before occupants are allowed to re-enter.

FEDERALLY INSPECTED MEAT & POULTRY ESTABLISHMENTS: Apply as a space or directed contact spray for control of flying and crawling insects listed (see "Insects controlled" section) in establishments operating under the Federal meat, poultry, shell egg grading and egg products inspection program. Do not apply while food processing is

underway. All food processing surfaces and equipment should be covered or washed with an effective cleaning compound followed by a potable water rinse after application and before processing resumes. Close doors, windows and shut off all ventilating equipment prior to application. When using ULV (Ultra Low Volume) equipment, adjust it to deliver aerosol droplets (30 microns or less) and apply 1 oz./1,000 ft.³ of space. Leave room closed for at least 1 hour after application and ventilate before occupants are allowed to re-enter.

DEEP VOID APPLICATION

SPOT APPLICATION: With equipment designed for deep void Ultra Low Volume (ULV) applications follow the manufacturer's recommendations for proper equipment set up and operating pressures. Place the tip of the applicator at or into the crack and crevice to be treated. Apply product in short bursts while keeping tip in position. Allow the air stream to push insecticide deep into voids. Repeat application as necessary to flush insects from voids. Avoid applying to the point of runoff or drip. Ventilate thoroughly before occupants are allowed to re-enter.

LINEAR APPLICATION ALONG BASEBOARDS, TRIM MOLDING, ETC.: With equipment designed for deep void ULV application, move applicator tip along the crevice to be treated applying a constant burst of insecticide. Move the tip at a steady rate of approximately 1 ft./second. Avoid applying to the point of runoff or drip. Repeat application as necessary to flush insects from the void. Ventilate thoroughly before occupants are allowed to re-enter.

SPOT, OR CRACK AND CREVICE APPLICATIONS - UNDILUTED: Use concentrate undiluted and apply through equipment which can deliver a pin stream spray. Apply in specific areas such as: cracks and crevices in or behind baseboards, in floors, walls, expansion joints, areas around water and sewer pipes, wall voids, or voids in table legs and equipment where pests can hide as well as other similar areas. Spot treatments may also be made to areas including but not limited to storage areas, closets, doors and windows, behind and under refrigerators, cabinets, sinks, stoves, and other equipment, the underside of shelves, drawers and similar areas. Do not apply where electrical short circuits could occur.

DILUTED SPRAY APPLICATIONS

SURFACE SPRAY (INCLUDING SPOT, CRACK AND CREVICE TREATMENTS):

Use diluted concentrate with sprayers that give large, wetting spray particles, to insure maximum control of listed crawling insects (see "Insects Controlled" section): Dilute 1 part concentrate with up to 9 parts oil and mix well. Apply as a coarse wetting spray using approximately 1 gallon of spray mixture for 750 square feet of surface. Spray walls, floor and other surfaces thoroughly, being especially careful to apply the spray into all cracks and crevices in woodwork, walls, floors, underneath sinks, behind pipes and in all places where these insects shelter. Treat both the underside and topside of infested carpets. Spray inside of trunks, closets and infested areas of shelving. Treat unloading, storage, handling and processing areas, in and around bins, conveying, processing and handling equipment. Spray thoroughly around the base of heavy machinery and equipment. Apply around doors and windows, behind and under refrigerators, cabinets, sinks, stoves, and other equipment, the underside of shelves, drawers and similar areas. Do not apply where electrical short circuits could occur. Spray insects directly when possible.

TO KILL ACCESSIBLE STAGES OF GRANARY INSECTS: Dilute 1 part Concentrate with up to 9 parts of oil and mix well. Spray outside walls of grain bins, flooring, about base of milling machinery, walls and baseboards, areas behind cabinets, around pipes and other insect harborages using 1 gallon spray per 1,000 square feet of surface.

ANIMAL HOUSING AND PRODUCTION FACILITIES: Use to control Flying and Crawling Insects listed in the "Insects Controlled" section of this label. Apply as a Space Spray application, Crack and Crevice or Spot Treatment, or as a diluted surface spray. Animals raised for milk and meat production must not be present at the time of application. They may be returned to the facility following ventilation (Space Spray) or once the sprayed material has dried (Surface Application).

SPACE SPRAY APPLICATION: To control listed crawling and flying insects. Close doors, windows and shut off all ventilating equipment. Then with a mechanical aerosol generator (no droplets over 50 microns in diameter and 80% less than 30 microns) apply undiluted concentrate at a dosage rate of approximately 1 fl. oz./1,000 ft.³ of room space. Direct spray toward areas where insects travel, land, harbor and breed, including walls, cages, shelves, ceiling, rafters, support beams, posts, upper areas of facility. Do not allow application to contact bulk animal feedstuffs or drinking water. Do not remain in treated areas after application, and leave facility closed for at least 1 hour. Ventilate thoroughly before milk and meat production animals are allowed to re-enter.

SURFACE SPRAY APPLICATION (INCLUDING SPOT, CRACK AND CREVICE

TREATMENTS): Use diluted concentrate with sprayers that give large, wetting spray particles, to insure maximum control of listed insects (see "Insects Controlled" section): Dilute 1 part concentrate with up to 9 parts oil and mix well. Apply as a coarse wetting spray using approximately 1 gallon of spray mixture for 1,000 square feet of surface. Spray walls, floor and other surfaces thoroughly, being especially careful to apply the spray into all cracks and crevices, on walls, floors, ceilings, around windows and doors, behind pipes and in all places where these insects shelter. Treat exterior surface of bins, conveying, and handling equipment. Do not apply where electrical short circuits could occur. Spray insects directly when possible. Animals may be returned to treated area once applied spray has dried.

For best results, use as part of an integrated program with a residual spray such as Onslaught® Microencapsulated Insecticide, and with an Insect Growth Regulator such as NyGuard® IGR Concentrate.

RESTRICTIONS

Product cannot be used: 1) in areas where animals raised for milk and meat (human food animals) are present; 2) in any areas where animal feeds are prepared, processed, or stored, and; 3) in any truck beds or truck trailers, freight/tank/container, railroad cars, ship holds, or ship and air freight containers that are used to transport unbagged (bulk) feeds.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry area. Always store pesticides in the original container. Store away from food and pet food.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

For containers with capacities equal to or less than 5 Gallons (18.9 L):

Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Contains petroleum distillates. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt and long pants, socks, shoes, and gloves. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Wear a respirator with an organic-vapor removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any R, P or HE prefilter when applying this product as a space spray.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF SWALLOWED:

- Immediately call a poison control center or doctor.
- Do not induce vomiting unless told to do so by a poison control center or a doctor.
- Do not give any liquid to the person.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance. Then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Contains petroleum distillate – vomiting may cause aspiration pneumonia. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-888-740-8712.

alto**CIRRUS**[®]
Fogging Concentrate

EPA Reg. No. 1021-1875
EPA Est. No. 1021-MN-2
0510-0911



ETOC is a registered trademark of Sumitomo Chemical Corp., Ltd., Japan.
All other marks are property of their respective holders. Manufactured by MGK[®].
©2011 MGK. All rights reserved.
8810 Tenth Avenue North, Minneapolis, MN 55427

2.5M-1111